



Literature Review on the Welfare Implications of
Tail Docking of Cattle

(August 29, 2014)

THE ISSUE

Tail docking is a management practice used within both the dairy and beef industries. The dairy industry in New Zealand developed the process during the early 1900s as an attempt to reduce the incidence of leptospirosis in milking personnel. The stated goals of tail docking in dairy cows include improved comfort for milking personnel, enhanced udder cleanliness, reduced incidence of mastitis, and improved milk quality and milk hygiene. For beef cattle, tail docking is used in confined slatted floor feedlot operations; these facilities are mainly located in the Great Lakes region of the United States and Ontario Canada within North America. Stated goals are to reduce injury associated with tails being stepped on by other cattle and/or caught in between the slats of slatted floors, and to prevent subsequent tail infection, ascending myelitis, septicemia, and lameness resulting from these injuries.

A variety of methods have been used to dock tails in dairy cattle, including cauterizing docking irons, application of elastrator bands, use of emasculators, and surgical excision. The application of elastrator bands is the most commonly employed method. Tail docking in the dairy industry is usually performed on preparturient heifers or calves near weaning age. An elastrator band or tight rubber ring is applied to the tail so that between 1/3 and 2/3 of the tail are removed; in New Zealand, regulations determine the minimum length between the distal vulva and the site of band application.¹ Placement distal to the sixth coccygeal vertebra has been recommended to ease the docking process and to avoid leaving a tail that is too short for proper restraint or that parts the vulvar lips and allows manure contamination of the urogenital tract.² The necrotic distal portion of the tail detaches 3 to 7 weeks after banding, or may be removed by using clean shears. Tail docking in the beef industry at the feedlot requires that the distal 2/3 of the tail be removed immediately after placement of the elastrator band or rubber ring.

TAIL DOCKING IN THE UNITED STATES AND OTHER COUNTRIES

Tail docking is no longer a common procedure in dairies in New Zealand, and the practice appears to be declining in other countries including the United States, although it remains common in some geographic regions. Denmark, Germany, Scotland, Sweden, the United Kingdom, and some Australian states prohibit tail docking. In Australian states where the practice is permitted, guidelines state it should be performed when recommended by a veterinarian for health reasons, and the tail stump must be long enough to cover the vulva.² In Canada, national guidelines recommend that the procedure be performed on young calves by trained personnel with the proper equipment and attention to pain relief.² The Canadian Veterinary Medical Association opposes the practice of docking of dairy cattle for management purposes.³

In the United States, California has passed legislation banning routine tail docking in dairy cattle and similar actions have been proposed in other states. A survey of 113 North Central and North Eastern U.S. dairies⁴ found that tail docking was practiced on 82.3% of the dairies. The most common reported

This peer-reviewed summary has been prepared by the American Veterinary Medical Association Animal Welfare Division. While principally a review of the scientific literature, it may also include information gleaned from proprietary data, legislative and regulatory review, market conditions, and scholarly ethical assessments. It is provided as information and its contents should not be construed as official AVMA policy. Mention of trade names, products, commercial practices or organizations does not imply endorsement by the American Veterinary Medical Association.

© American Veterinary Medical Association

docking time in dairy cattle was before or shortly after calving (35.2%). Rubber band was the most common method (92.5%) in dairies. Cow hygiene was suggested as the most common reason to dock (73.5%) dairy cows followed by worker comfort at 17.4%.

Current AVMA policy opposes routine tail docking of cattle. The current position of the American Association of Bovine Practitioners (AABP) states that *“The AABP opposes the routine tail docking of cattle. Current scientific literature indicates that routine tail docking provides no benefit to the animal.”*⁵

SCIENTIFIC EVIDENCE ON THE BENEFITS OF TAIL DOCKING

Reduced risk of leptospirosis in milkers—Urine from infected animals is the primary source of transmission of leptospirosis. Infection can occur via contact with skin abrasions or wounds, or via contact with the mucous membranes of the eyes, nose, and mouth.⁶ Docking is thought to reduce the risk of leptospirosis by eliminating the possibility that a urine-soaked tail could contact the milker’s skin or face. Mackintosh et al⁷ determined that the leptospiral titers of milkers bore no relationship to tail docking, and hypothesized that transmission of leptospirosis in endemic herds likely occurs from sources other than tail contact.

Improved cow and udder cleanliness—Anecdotal support for tail docking centers on the idea that a soiled tail can inoculate the udder with pathogens. One study revealed that rear-quarter cleanliness was greater for docked cows compared with intact cows; however, no statistical differences were observed with respect to udder cleanliness or somatic cell count (SCC).⁸ In another study, cow cleanliness, udder cleanliness, and SCC scores were not different for docked heifers compared with intact heifers.⁹

Reduced incidence of mastitis and improved milk hygiene—Environmental pathogens present in dirt, manure, and water can cause mastitis in dairy cattle. Tail docking is reported to decrease the incidence of mastitis caused by environmental pathogens by eliminating the possibility that a heavily soiled tail or tail switch would come in contact with the udder. A review of the related scientific literature reveals leg cleanliness scores were improved in docked cattle compared with intact cattle. No significant differences were observed in SCC, udder cleanliness, or intramammary infection between docked and intact cattle.¹⁰ Although docked cattle had a higher incidence of mastitis in one study,⁹ the difference was not statistically significant.

Reduced incidence tail injury and improved performance—Trampling by pen mates appears to be the major cause of tail damage in indoor beef feedlots. The tail tip of the lying animal lies on the floor away from the animal’s body and is unprotected against trampling. A tail that is lying on a hard, inflexible sharp-edged surface (e.g., a slatted floor) is more likely to incur severe damage from trampling than a tail that is on a soft, flexible surface. In general, slatted floor facilities have higher stocking densities than those with solid floors, and slatted facilities with the highest stocking densities had the highest prevalence of tail tip necrosis.^{11,12}

Tail tip necrosis can lead to tail infection, ascending myelitis, septicemia, and lameness in cattle that have suffered a tail injury however the risk of subsequent infection from tail injury is low. A study on the prevalence of tail tip necrosis in Ontario Canada found 34.5% of tails inspected at slaughter plants to have necrosis with only 3.4% having infections.¹³ Another study conducted in Ontario found that cattle housed in solid floor facilities had no tail tip necrosis, whereas 1.36% of cattle housed in slatted floor facilities were treated or slaughtered for tail tip necrosis.¹¹ A similar study in Nebraska found 1% of cattle housed in a slatted floor facility to have tail tip necrosis.¹⁴ Tail tip lesions occur most often in cattle without docked tails housed on slats, followed by cattle with docked tails housed on slats.¹² The lowest prevalence of tail tip lesions is in cattle housed in solid bedded facilities.¹¹ Severe tail tip lesions occur the most in cattle that are not tail docked and are housed on slatted floors.¹¹ Severe lesions also

occur in docked cattle housed on slats however no severe lesions were found on cattle housed in solid bedded facilities.¹¹ Finally, two studies identified no differences in performance or health indices between cattle with docked tails and cattle without docked tails housed on slats.^{15,16}

WELFARE CONCERNS—THE SCIENCE, RISKS, SEVERITY, AND ADVANTAGES

Acute pain—Application of a tight band around the tail produces extraluminal compression of the arteries and veins, resulting in impeded arterial flow and venous drainage in the tissues. Lack of perfusion compromises the supply of oxygen and metabolic substrates to the tissues and results in ischemia. Continued ischemia induces severe cellular damage and coagulation necrosis.^{17,18} Ischemic lesions of the intestinal tract or limbs are known to cause pain during the acute phase, followed by decreased pain as the lesion progresses. The greatest challenge for determining the severity of pain associated with tail docking in cattle lies in an accurate assessment of signs of pain in this species.

Three-week-old Holstein calves have been observed to exhibit increased walking or running behavior, increased head-to-tail movement and licking, and less tail swinging and lying behavior following application of a rubber ring for tail docking. These actions have been interpreted as indicators of mild distress. Distal tail sensitivity to hot water was absent 75 to 105 minutes after banding, indicating desensitization of the tail below the banding site.¹⁹

When compared with 7- to 21-day-old calves, banded 22- to 42-day-old calves exhibited significantly more rear visualization and restlessness than intact calves of the same age.²⁰ That study's investigators concluded that, although age-related behavioral differences were observed, tail docking of calves produced a minor response.

Based on measurement of plasma cortisol concentrations of calves before and after tail docking with a rubber ring or cauterizing iron, Petrie et al²¹ determined that docking did not result in significantly more distress than restraint and blood sampling. A small number of 3- to 4-month-old calves exhibited tail shaking and vocalization, which was interpreted as discomfort. The use of local anesthetic at the time of ring application provided no detectable benefit in reducing physiologic signs of stress.

Tail docking of adult heifers using an elastrator band with or without local anesthesia resulted in few physiologic or behavioral effects. Banded heifers spent more time eating during the week following banding, which may have represented displacement behavior and mild distress; these behaviors returned to pre-banding levels when necrotic tails were removed. Increases in plasma cortisol concentrations, considered indicators of stress, were not observed.²² Restlessness was observed during the first hour after banding in preparturient heifers, but the finding was not statistically significant.¹⁰

Docked heifers spent significantly less time holding their tails in the raised position following docking and significantly more time with their tails in the pressed position, but no differences were observed in feed intake or milk production. Holding the tail in the pressed position was interpreted as an indicator of discomfort. Significantly less tail shaking was observed in docked heifers, and was hypothesized to be related to pain associated with shaking the recently banded tail. No differences were observed in behavior of heifers docked with or without epidural anesthesia. The investigators involved in that study concluded that tail docking was associated with minimal discomfort in heifers, and that use of epidural anesthesia provided no benefit.²³

Chronic pain—Following trauma to peripheral nerves (including that induced by banding and docking), continued growth of damaged nerve axons may result in the formation of a mass of tangled axons called a neuroma.²⁴ Neuromas are associated with chronic pain, and may play a role in post-amputation pain in humans.² Neuromas have been reported to develop after beak trimming in poultry

and tail docking in pigs,^{25,26} and were observed at slaughter in tail stumps of adult cattle that had been docked using a knife at 12 to 18 months of age.²⁷ Eicher et al²⁸ documented increased sensitivity to heat or cold in previously docked heifers. These findings were comparable to those observed in humans experiencing phantom limb pain following amputation, and were interpreted as indicators of chronic pain. In addition, neuromas were identified in the tail stumps of the docked heifers included in the study.

Physiologic stress—Blood cortisol concentrations have been studied as indicators of physiologic stress in animals. Tail docking of preparturient Holstein heifers did not result in significant alterations in cortisol concentrations.^{8,20,22} Docking of three-week-old calves did not significantly increase blood cortisol concentrations above those associated with handling and sample collection.²¹

Disease—Necrotic tissue, such as the ischemic distal tail after banding, is prone to infection with pathogens. Clostridial organisms, ubiquitous in soil, may colonize the wound and result in local or systemic infection. Tetanus and gangrene have been reported after tail docking, and vaccination against clostridia is recommended prior to performing the procedure.²

Behavior—The role of the tail in communication between cattle has not been documented, but it has been speculated that tail docking limits the ability of cattle to exhibit normal signaling behavior.^{29,30} In addition, the tail is widely believed to play a role in fly control; shaking the tail and brushing the body and limbs can dislodge biting flies. Fly avoidance behaviors following tail docking have received attention from researchers for their implications for animal welfare. The stable fly (*Stomoxys calcitrans*) is a common disruptive fly, and its presence has been associated with increased stress, reduced milk production and weight gain, disrupted grazing, and reduced growth.³¹ Observed fly avoidance behaviors include stomping, kicking the trunk, tail swishing, skin twitching (panniculus reflex), head and ear motion, and taking flight.^{8,19}

Fly counts have been observed to be greater on the rear limbs of docked three-week-old calves during times of high fly activity.¹⁹ Several studies have confirmed that although front limb fly avoidance behaviors did not differ, rear limb fly avoidance behaviors were significantly increased in docked cows when compared with intact cows.^{8,32,33,34} In addition, almost twice as many flies were observed on the rear limbs of docked cows compared with control cows.⁷

SUMMARY

Anecdotal reports of the benefits of tail docking are not currently supported by data in the scientific literature. Tail injury from trampling can be minimized by maintaining a lower stocking density and providing solid flooring and/or bedding for cattle. Tail docking has been experimentally shown to cause minimal adverse physiologic effects; however, fly avoidance behaviors are more frequent in docked cattle, suggesting potential long-term adverse behavioral effects. Increased temperature sensitivity and the presence of neuromas suggest that chronic pain may be associated with the procedure.

REFERENCES

1. New Zealand Ministry of Agriculture and Forestry website. Available at: <http://www.biosecurity.govt.nz/animal-welfare/codes/dairy-cattle/index.htm>
2. Stull CL, Payne MA, Berry SL, et al. Evaluation of the scientific justification for tail docking in dairy cattle. *J Amer Vet Med Assoc* 2002; 220(9): 1298-1302.
3. Canadian Veterinary Medical Association. Available at: <http://canadianveterinarians.net/ShowText.aspx?ResourceID=25>
4. Fulwider WK, Grandin T, Rollin BE et al. Survey of dairy management on one hundred thirteen north central and northeastern United States dairies. *J Dairy Sci* 2008; 91: 1686-1692.

5. American Association of Bovine Practitioners current position on tail docking in cattle, approved by AABP Board, March 13, 2010. American Association of Bovine Practitioners, P.O. Box 3610, Auburn, AL 36831. <http://www.aabp.org>
6. Government of Western Australia Department of Agriculture Farmnote, no.14/2003: Leptospirosis. Available at: www.agric.wa.gov.au
7. Mackintosh CG, Schollum LM, Blackmore DK, et al. Epidemiology of leptospirosis in dairy farm workers in the Manawatu, part II: a case-control study of high and low risk farms. *NZ Vet J* 1982; 30: 73-76.
8. Eicher SD, Morrow-Tesch, JL, Albright JL, et al. Tail-docking alters fly numbers, fly-avoidance behaviors, and cleanliness, but not physiological measures. *J Dairy Sci* 2001; 84: 1822-1828.
9. Tucker CB, Fraser D, Weary DM. Tail docking dairy cattle: effects on cow cleanliness and udder health. *J Dairy Sci* 2001; 84: 84-87.
10. Schreiner DA, Ruegg PL. Effects of tail docking on milk quality and cow cleanliness. *J Dairy Sci* 2002; 85: 2503-2511.
- 11 Drolia H, Luescher UA and Meek AH. Tail-tip necrosis in Ontario feedlot cattle: two case-control studies. *Prev Vet Med.* 1990;9:195-205
- 12 Schrader L, Roth H-R, Winterling C, et al. The occurrence of tail tip alteration in fattening bulls kept under different husbandry conditions. *Ani Welfare.* 2001;10:119-130.
- 13 Drolia H, Luescher UA, Meek AH et al. Tail tip necrosis in Ontario beef feedlot cattle. *Can Vet J.* 1991;32:23-29.
- 14 Thomson DU, Taylor W, Noffsinger T, et al. Case report-tail tip necrosis in a confined cattle feeding operation. *Bovine Pract.* 2009;43(1):18-22.
- 15 Grooms D, Schwehofer J, Metz K, et al. Effect of tail docking on health and performance on feedlot calves housed in confined slotted floor facilities. *AABP Proceedings.* 2010;43:236.
16. Kroll, LK, Grooms DL, Siegford JM et al. Effects of tail docking on health and performance of beef cattle in confined, slatted floor feedlots. *J Anim Sci.* 2014; Jul 1[Epub ahead of print].
17. Kumar V, Abbas AK, Fausto N. Cellular adaptations, cell injury, and cell death. In: Kumar V, Abbas AK, Fausto N. Robbins and Cotran Pathologic Basis of Disease. 7th ed. Philadelphia: Elsevier, 2005; 3-46.
18. Mitchell RN. Hemodynamic disorders, thromboembolic disease, and shock. In: Robbins and Cotran Pathologic Basis of Disease. 7th ed. Philadelphia: Elsevier, 2005; 119-144.
19. Eicher SD, Dailey JW. Indicators of acute pain and fly avoidance behaviors in Holstein calves following tail-docking. *J Dairy Sci* 2002; 85: 2850-2858.
20. Schreiner DA, Ruegg PL. Responses to tail docking in calves and heifers. *J Dairy Sci* 2002; 85: 3287-3296.
21. Petrie NJ, Mellor DJ, Stafford KJ, Bruce RA, et al. Cortisol responses of calves to two methods of tail docking used with or without local anaesthetic. *NZ Vet J* 1996; 44(1): 4-8.
22. Eicher SD, Morrow-Tesch JL, Albright JL, et al. Tail-docking influences on behavioral, immunological, and endocrine responses in dairy heifers. *J Dairy Sci* 2000; 83: 1456-1462.
23. Tom EM, Duncan IJH, Widowski TM, et al. Effects of tail docking using a rubber ring with or without anesthetic on behavior and production of lactating cows. *J Dairy Sci* 2002; 85: 2257-2265.
24. Anthony DC, Frosch MP, De Girolami U. Peripheral nerve and skeletal muscle. In: Robbins and Cotran Pathologic Basis of Disease. 7th ed. Philadelphia: Elsevier, 2005; 1325-1347.
25. Breward J, Gentle MJ. Neuroma formation and abnormal afferent nerve discharges after partial beak amputation (beak trimming) in poultry. *Experientia* 1985; 41: 1132-1134.
26. Simonsen HB, Klinken L, Bindseil E. Histopathology of intact and docked pigtailed. *Br Vet J* 1991; 147: 407-412.
27. Barnett JL, Coleman GJ, Hemsworth PH, et al. Tail docking and beliefs about the practice in the Victorian dairy industry. *Aust Vet J* 1999; 77(11): 742-747.
28. Eicher SD, Cheng HW, Sorrells AD, et al. Short communications: behavioral and physiological indicators of sensitivity or chronic pain following tail-docking. Accepted, *J Dairy Sci.* 2005; Manuscript on file.
29. Kahler SC. Studies offer scientific appraisal of tail docking effects in dairy cattle. *J Amer Vet Med Assoc* 2002; 221(12): 1674.
30. American Veterinary Medical Association. DiVita LJ. Natural bovine behavior key to evaluating management practices. Available at: <http://www.avma.org/onlnews/javma/jan01/s011501pp.asp>
31. Campbell JB, Berry IL. Economic threshold for stable flies on confined livestock. *Misc Publ Entomol Soc Am No* 1989; 74: 18-22.
32. Phipps AM, Mathews LR, Verkerk GA. Tail docked dairy cattle: fly induced behaviour and adrenal responsiveness to ACTH. *Proc NZ Soc Anim Prod* 1995; 55: 61-63.
33. Mathews LR, Phipps RA, Verkerk D, et al. The effects of taildocking and trimming on milker comfort and dairy cattle health, welfare and production. *Anim Beh Welf Res Cent* 1995, 1-25.
34. Wilson GDA. Docking cows' tails. *Ruakura Farm Conf (Ruakura, NZ)* 1972; 158-165.