

PROHIBIT[®] Soluble Drench Powder

(Levamisole Hydrochloride)



Prohibit Soluble Powder is a broad spectrum anthelmintic effective against stomach, intestinal and lungworms in cattle and sheep.

- Available in 52 gram pouch and 605 gram bottle
- Broad coverage of nine major types of worms
- Short withdrawal time
- Cost-effective and well-targeted
- For oral use in cattle and sheep

Available in the following presentations:

- (30 x 52 gram pouch bucket)
- (15 x 52 gram pouch per display box)
- (605 gram soluble drench powder)

DIRECTIONS AND DOSAGES

Cattle and sheep de-wormer. For oral use.

CATTLE STANDARD DRENCH SOLUTION - Place the contents of each pouch in a 1 quart (32 fl. oz.) container, fill with water; swirl until dissolved. Administer as a single drench dose according to the table on reverse side.

SHEEP STANDARD DRENCH SOLUTION - Place the contents of each pouch in a one gallon (128 fl. oz.) container, fill with water; swirl until dissolved. Administer as a single drench dose according to the table on reverse side.

Read full label and directions before use.



PROHIBIT[®] Soluble Drench Powder



Levamisole Hydrochloride Anthelmintic

Each 52 g pouch contains 46.8 g of levamisole hydrochloride activity.

ANADA 200-225, Approved by FDA

Each 605 g bottle contains 544.5 g of levamisole hydrochloride activity.

ANADA 200-225, Approved by FDA

Cattle and Sheep Dewormer. For Oral Use.

Administer as a standard drench with standard drench syringe or administer as a concentrated drench solution with an automatic drenching syringe.

Mixes as a clear solution.

INDICATIONS:

Prohibit[®] (levamisole hydrochloride) is a broad-spectrum anthelmintic and is effective against the following nematode infections in cattle and sheep:

STOMACH WORMS: (*Haemonchus*, *Trichostrongylus*, *Ostertagia*)

INTESTINAL WORMS: (*Trichostrongylus*, *Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*) [*Chabertia* - sheep only]

LUNGWORMS: (*Dictyocaulus*)

DOSAGE AND ADMINISTRATION

CATTLE-STANDARD DRENCH SOLUTION- Place the contents of each packet in a 1 quart (32 fl. oz.) container, fill with water; swirl until dissolved. Administer as a single drench dose according to the following table:

Weight	Drench Dosage	Each Pouch Will Treat	CATTLE
200 lb	1/2 fl oz	64 head	
400 lb	1 fl oz	32 head	
600 lb	1 1/2 fl oz	21 head	
800 lb	2 fl oz	16 head	

CATTLE-CONCENTRATED DRENCH SOLUTION- For use with automatic syringe. Place the contents of each pouch in a standard household measuring container and add water to the 8 3/4 fl oz level. Swirl until dissolved. Give 2 mL (milliliter) per 100 lb body weight. Refer to the table above for the number of cattle each packet will treat.

SHEEP-STANDARD DRENCH SOLUTION- Place the contents of each packet in a 1 gallon (128 fl oz) container, fill with water, swirl until dissolved. Administer as a single drench dose according to the following table:

Weight	Drench Dosage	Each Pouch Will Treat	SHEEP
50 lb	1/2 fl oz	256 head	
100 lb	1 fl oz	128 head	
150 lb	1 1/2 fl oz	84 head	
200 lb	2 fl oz	64 head	

SHEEP-CONCENTRATED DRENCH SOLUTION- For use with automatic syringe. Place the contents of each packet in a standard household measuring container and add water to the 17 1/2 fl. oz. level. Swirl until dissolved. Give 2 mL per 50 lb. body weight. Refer to the previous table for the number of sheep each packet will treat.

GALLON BOTTLE - When you are ready to de-worm your cattle or sheep, add water to the powder in this bottle up to the 3 liter mark. Swirl to mix thoroughly before using. If any solution is left over, it may be stored for up to 3 months in this tightly capped bottle, shake well before using.

Administer as a single drench dose as follows:

Cattle -- 2 mL per 100 lb. body weight

Weight	Drench Dosage	Each Bottle Will Treat	CATTLE
100 lb	2 mL	1,500 head	
300 lb	6 mL	500 head	
500 lb	10 mL	300 head	
700 lb	14 mL	214 head	
1,000 lb.	20 mL	150 head	

Sheep -- 1 mL per 50 lb. body weight

Weight	Drench Dosage	Each Bottle Will Treat	SHEEP
50 lb	1 mL	3,000 head	
100 lb	2 mL	1,500 head	
150 lb	3 mL	1,000 head	
200 lb	4 mL	750 head	

NOTE: Careful weight estimates are essential for proper performance of this product. Prepare solutions as needed. However, excess solutions may be stored in clean closed containers up to 90 days without loss of anthelmintic activity.

Cattle and sheep maintained under conditions of constant helminth exposure may require retreatment within 2 to 4 weeks after the first treatment.

WARNING: Do not administer to cattle within 48 hours of slaughter for food. Do not administer to sheep within 72 hours of slaughter for food. To prevent residues in milk, do not administer to dairy animals of breeding age.

CAUTION: Muzzle foam may be observed. However, this reaction will disappear within a few hours. If this condition persists, a veterinarian should be consulted. Follow recommended dosage carefully. Consult veterinarian before using in severely debilitated animals.

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

Store between 15° and 30°C (59° and 86°F). **WARNING:** Keep out of reach of children. For Animal Use Only

© Registered trademark of Huvepharma, Inc.

Manufactured for Huvepharma, Inc.

525 Westpark Drive, Suite 230, Peachtree City, GA 30269

www.huvepharma.us 877.994.4883

ANAD-0048 (Rev 1019)



Read actual product label
directions thoroughly before use.