

CATTLE VACCINE REFERENCE CHART

	LABEL CLAIMS						SCOURS		CLOSTRIDIAL	FOOTROT	PINKEYE		
	Approved for Pregnant Cows and Heifers	Withdrawal Period	Minimum Age Restrictions	Dose	Administration	Adjuvant	E. coli	Rotavirus	Coronavirus	C. perfringens Type A	C. perfringens Type C	Fusobacterium necrophorum	Moroxella bovis
Bovine Pili Shield®	■	60 days	—	1 mL	IM	Xtend III	■						
Bovine Pili Shield + C	■	60 days	—	1 mL	IM	Xtend III	■			■			
Scour Bos® 4	■	60 days	—	2 mL	IM	Xtend III		■	■				
Scour Bos 9	■	60 days	—	2 mL	IM	Xtend III	■	■	■	■			
Clostridium Perfringens Type A Toxoid		21 days	—	2 mL	SQ	Proprietary dual component			■				
Fusogard®		21 days	6 months	2 mL	SQ	SuprImm®				■			
Pinkeye Shield® XT4		60 days	—	2 mL	IM	Xtend SP					■		

Bovine Pili Shield®



Clostridium Perfringens Type A Toxoid

Fusogard®

Pinkeye Shield® XT4

To learn more about the portfolio of Elanco cattle vaccines, contact your herd health veterinarian, Elanco sales representative or technical consultant.

For all products: The label contains complete use information, including cautions and warnings.
Always read, understand and follow the label and use directions.